

INSPIRATORY RESISTANCE MAINTAINS ARTERIAL PRESSURE DURING CENTRAL HYPOVOLEMIA: IMPLICATIONS FOR TREATMENT OF COMBAT CASUALTIES WITH SEVERE HEMORRHAGE

Victor A. Convertino*, Kathy L. Ryan, Caroline A. Rickards, and John B. Holcomb
US Army Institute of Surgical Research, Fort Sam Houston, TX 78234

William H. Cooke
University of Texas at San Antonio, San Antonio, TX 78249

Ahamed H. Idris
Dept. of Surgery & Emergency Medicine, UT Southwestern Medical Center, Dallas, TX 75235

Anja Metzger and Keith G. Lurie
Advanced Circulatory Systems, Inc., Eden Prairie, MN 55344

Bruce D. Adams
Emergency Medicine Department, Brooke Army Medical Center, Fort Sam Houston, TX 78234

ABSTRACT

Loss of consciousness due to central hypovolemia can occur due to sudden cardiovascular decompensation in normal individuals or hypovolemic shock in wounded patients. A variety of devices have been developed to sustain perfusion to the brain including anti-G suits worn by pilots and returning astronauts and applied to patients as “shock trousers.” However, all countermeasures developed to date suffer from problems that limit their utility in the field. An “impedance threshold device” (ITD) has recently been developed that acutely increases central blood volume by forcing the thoracic muscles to develop increased negative pressure, thus drawing venous blood from extrathoracic cavities into the heart and lungs. We review here a series of experiments that demonstrate the application of the ITD to a variety of experimental conditions, including its use to: (a) increase heart rate, stroke volume, and arterial blood pressure in normovolemia and hypovolemia; (b) increase cerebral blood flow velocity; (c) reset cardiac baroreflex function to a higher operating range for blood pressure; (d) lower intracranial pressure; and (e) reduce orthostatic symptoms. In this brief review, we present evidence that supports further consideration of using inspiratory resistance as a countermeasure against circulatory collapse associated with orthostatic instability and hemorrhagic shock.

1. INTRODUCTION

Maintenance of consciousness requires adequate perfusion to the brain, which may be compromised in a

variety of physiological and clinical circumstances. Inability to tolerate upright standing posture due to development of severe orthostatic hypotension and syncope often plagues astronauts and military personnel in their austere operational environments. In the civilian sector, up to 30% of otherwise healthy young adults report at least one syncopal episode during their lifetimes, and syncope accounts for up to 3% of all emergency room visits in the United States (Kapoor, 1996). More critically, hemorrhagic shock remains a leading cause of death in both civilian and battlefield trauma (Carrico et al., 2002). Syncope and hemorrhagic shock share the same underlying mechanisms, namely, central hypovolemia and cardiovascular decompensation. A countermeasure that functionally restores central blood volume would therefore be expected to prove useful for all of these conditions.

Battlefield injury often leads to hypovolemia through hemorrhage. In spaceflight, hypovolemia occurs as a response to microgravity over the first several days of exposure (Convertino, 1996), persists regardless of flight duration, and contributes to post flight orthostatic intolerance and reduced exercise capacity (Convertino et al., 1996; Levine et al. 1996). The usual countermeasures for all of these conditions include fluid replacement (resuscitation) and/or lower body counter pressure (shock trousers or G-suits). For spaceflight these often fail to prevent symptoms of cardiovascular instability or even frank syncope upon assuming an upright body position at 1 G (Buckey et al. 1996). Tolerance for loss of central blood volume or orthostatic stress can be enhanced by means of centrifuge training (Evans et al., 2004) and maximal exercise bouts prior to orthostatic testing

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 01 NOV 2006		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE Inspiratory Resistance Maintains Arterial Pressure During Central Hypovolemia: Implications For Treatment Of Combat Casualties With Severe Hemorrhage				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) US Army Institute of Surgical Research, Fort Sam Houston, TX 78234				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADM002075., The original document contains color images.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 8	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

(Engelke et al., 1996), but these procedures cannot be applied in space craft and remote settings or medical evacuation aircraft.

Low central blood volume contributes to a reduction in cardiac filling, stroke volume (SV), and arterial pressure (AP). The resulting acute hypotension activates autonomically-mediated compensatory mechanisms that evoke sympathetic nerve activity, tachycardia and peripheral vasoconstriction in an attempt to restore AP (10). When the reduction in blood volume and AP reach a critical level, activation of decompensatory mechanisms result in sympathetic withdrawal, bradycardia and vasodilation (Convertino and Cooke, 2002); a condition we refer to as circulatory collapse (Convertino et al., 2004a; Cooke et al., 2004). Cardiovascular decompensation is the precursor to syncope or hemorrhagic shock. Therefore, any therapeutic approach that is designed to increase venous return and SV should counteract circulatory collapse. Increased negative intrathoracic pressure during spontaneous inspiration represents a natural mechanism for enhancing venous return and cardiac filling. Any device that applies resistance during inspiration takes advantage of this simple concept and shows promise as a mechanical facilitator of the respiratory pump that enhances venous return and preload to the heart (Lurie et al., 1995, 2004; Samniah et al., 2003). In this review, we examine a series of experiments designed to evaluate the application of inspiratory resistance as a potential countermeasure to restore central blood volume and possibly improve clinical and critical outcomes.

2. USING THE CHEST AS A VACUUM PUMP

Cournand et al. (1948) reported in an early study that venous return, ventricular preload, and subsequently cardiac output (CO) decreases with positive pressure breathing sufficient to increase mean airway pressure. Guyton et al. (1957) characterized the entire venous return curve in animals by plotting blood flow against pressures in the right atrium, and demonstrated marked increases in venous return when right atrial pressures were suctioned to -2 to -4 mmHg. Based on this simple concept of positive versus negative intrathoracic pressures and their effects on venous return, it has been shown that increasing negative intrathoracic pressure through resistive breathing decreases left ventricular and right atrial pressures (Lurie et al., 2004), consequently increasing left ventricular preload and SV index (Marino et al., 2004). The central hemodynamic response of resistive breathing is similar to that observed during Mueller maneuvers, where initial reductions of arterial pressure (AP) due to reductions of left ventricular stroke volumes are followed by increases in AP due to resulting increased venous return and consequent increases in left

ventricular SV (Fitzgerald et al., 1991; Morgan et al., 1993).

Negative intrathoracic pressure during the inspiration may be enhanced in several ways. In the experiments described here, a controlled level of inspiratory negative pressure was produced in humans by using an inspiratory threshold device (ITD) comprised of a plastic valve attached to a standard clinical facemask (Lurie et al., 2000). Responses to ITD breathing were compared directly with a sham ITD device which provided zero inspiratory pressure (ZTD).

3. CENTRAL HEMODYNAMICS

Changes of central hemodynamics in humans during resistive breathing were assessed in two human studies (Convertino et al., 2004b, 2004c). During spontaneous breathing in the supine position, inspiratory impedance of approximately 6 cm H₂O increased heart rate (HR), AP (Convertino et al., 2004b, 2004c), SV (measured with thoracic bioimpedance) and CO, and decreased total peripheral resistance (TPR) (Convertino et al., 2004c). Other countermeasures that restore central blood volume and protect SV and CO such as maximal exercise, G-suits, fluid loading, and centrifuge training may fall short of effective implementation due to practical limitations (Bungo et al., 1985; Engelke et al., 1996; Evans et al., 2004) or the inability to produce the immediate effects on central blood volume and hemodynamics similar to resistive breathing (Evans et al., 2004).

Although loss of blood volume contributes to postflight orthostatic intolerance in astronauts, reduction in the sensitivity of the carotid-cardiac baroreflex has also been implicated in hemodynamic instability following both simulated (Convertino et al., 1990; Eckberg et al., 1992) and actual microgravity (Fritsch-Yelle et al., 1994; Fritsch et al., 1992). Carotid-cardiac baroreflex function is restored in subjects after bedrest with application of maximal exercise 24 hours prior to reambulation (Engelke et al., 1996). Similarly, reductions of carotid-cardiac baroreflex sensitivity associated with reductions of central blood volume during lower body negative pressure (LBNP) are reversed with restoration of central volume during G-suit inflation (Eiken et al., 1991). Stimulation of arterial and/or cardiopulmonary baroreceptors by oscillations in intrathoracic or arterial pressure (Fitzgerald et al., 1981) may acutely change the sensitivity of the carotid-cardiac baroreflex response (Chapleau et al., 1987, 1989) and affect autonomic compensation to orthostatic or hypovolemic challenges.

4. AUTONOMIC FUNCTION

The effects of inspiratory resistance on the carotid-cardiac baroreflex response were also tested in humans. During ITD breathing, cardiac baroreflex sensitivity was not altered but responses were shifted to higher arterial pressures (Convertino et al., 2004b). These results, together with prior work (14), support the hypothesis that negative intrathoracic pressure and baroreflex resetting induced by ITD breathing augments central hemodynamics and potentially increases the operational range of the baroreflex under conditions of severe hypotension.

Increased HR in conjunction with increased AP could manifest through atrial stretch and activation of cardiopulmonary baroreceptors, but cardiopulmonary baroreflex activation and consequent interaction with arterial baroreceptors during inspiratory resistance can only be inferred and not measured directly in humans; it is likely that both cardiopulmonary and arterial baroreflexes operate to some degree and probably function at times in opposition (Barbieri et al., 2002). In addition, elevated HR and CO during spontaneous breathing on an ITD may simply reflect an “exercise” effect from the increased work of breathing against resistance. If this were true, one might expect withdrawal of vagal activity and no change or a slight increase in sympathetic activity with HR below 100 bpm (Rowell and Leary, 1990). However, in experiments designed to test the mechanism(s) involved in the tachycardic response to inspiratory resistance, there was no change in ventilatory mechanics (volume and rate), metabolic rate, cardiac vagal activity as indicated by no effect on the percent of normal consecutive R-R intervals that vary by more than 50 ms (pNN50) and muscle sympathetic nerve activity (microneurography) (Convertino et al., 2004b). While that report involved only one subject, later work on an additional eight subjects confirmed that ITD breathing does not affect vagal-cardiac control as estimated from frequency domain analysis of R-R intervals, or directly-measured peripheral sympathetic traffic (Cooke et al., 2006). Those observations suggest that the elevation in HR is initiated by a mechanical rather than metabolic or primary autonomic stimulus, and therefore may not represent an “exercise” effect per se. Rather, a larger negative intrathoracic pressure resulting from inspiratory resistance may initiate mechanically a chronotropic response as a result of enhanced cardiac filling [e.g., the Bainbridge reflex, stretch of the SA node (Bainbridge, 1915; Barbieri et al., 2002; Pawelczyk and Levine, 1995)].

Head-up tilt table experiments in astronauts prior to and immediately after the NASA Neurolab Space Mission (STS-90) revealed that increased muscle sympathetic nerve activity (MSNA) induced by moving from the supine to upright posture was associated with a reduction

in SV (Levine et al., 2002). Although this finding was not unexpected, lower average SV and greater average MSNA measured after space flight in both supine and upright postures were positioned in a linear fashion on the same SV-MSNA stimulus-response relationship as the average pre-flight SV and MSNA responses (Levine et al., 2002). Using LBNP as a model for the investigation of mechanisms associated with hemorrhagic shock (Cooke et al., 2004), we corroborated the linear relationship between SV and MSNA (Convertino and Cooke, 2002; Convertino et al., 2004a).

In addition to increasing cardiac filling and SV (Convertino et al., 2004c), spontaneous inspiration on the ITD lowered TPR (Convertino et al., 2004c). Since higher SV and lower TPR are associated with lower MSNA in a linear fashion (Convertino and Cooke, 2002; Convertino et al., 2004a; Levine et al., 2002), it seemed possible that spontaneous breathing on an ITD would cause a reduction in MSNA. However, recent experiments show that resistive breathing had no effect on supine MSNA (15 ± 8 vs. 15 ± 9 bursts/min) despite significant increases in mean arterial pressure (MAP) (94 ± 7 to 99 ± 9 mmHg) in eight normovolemic, normotensive subjects (Cooke et al., 2006). However, in one subject, a 23-ml (25%) increase in SV (measured with thoracic bioimpedance) during ITD breathing was associated with an MSNA of 23 bursts/min compared to 30 bursts/min when breathing on the ZTD (Fig. 1).




Fig. 1. Recording of neurograms from a subject during spontaneous breathing on a ZTD without inspiratory resistance (upper panel) and on an ITD (bottom panel). Figure from Convertino et al. [2004b].

Those preliminary results support the hypothesis that large elevations in SV might produce proportionate reductions in MSNA. Morgan et al. (1993) recorded MSNA responses within the respiratory cycle during prolonged (20-s) Mueller maneuvers and documented a

biphasic response consisting of initial suppression of sympathetic traffic despite falling AP followed by activation and resultant increases of AP. The ITD study averaged MSNA over several minutes; biphasic responses could have contributed to the observation there of unchanged sympathetic traffic during resistive breathing. Since high sympathetic nerve activity is associated with poor clinical outcome in states of central hypovolemia (Kleiger et al., 1987), the ability of the ITD to reduce MSNA could be an effective countermeasure against syncope and hemorrhagic shock. Future experiments designed to induce more dramatic alterations in central hemodynamics, particularly in states of central hypovolemia, are necessary to test this hypothesis.

5. CEREBRAL BLOOD FLOW

In a porcine model of cardiac arrest, cerebral blood flow (CBF) and neurological function were significantly protected by application of an ITD (Lurie et al., 1995, 2002). Yannopoulos et al. (in press) demonstrated in pigs that ITD breathing increases cerebral perfusion pressure (CPP), and increases CBF during cardiopulmonary resuscitation after cardiac arrest. Figure 2 shows a representative example of the changes in intrathoracic pressure measured in the trachea of the pig, and concurrent changes in intracranial pressure (ICP) measured in the brain parenchyma (Convertino et al., 2005a).




Figure 2. Continuous recordings of intrathoracic pressure (ITP, upper tracing) and intracranial pressure (ICP, lower tracing) in an apneic pig in hemorrhagic shock before, during (ITD on), and at the cessation (ITD off) of application of an ITD modified for use in nonbreathing individuals. Positive pressure ventilations were delivered every 8 seconds, when the ITD was applied. ITP was maintained at -10 mmHg in the absence of positive pressure ventilations. Figure from Convertino et al. [2005a]

In this case, positive pressure ventilations were delivered every 8 s and after each breath. Use of an ITD in conjunction with positive pressure breathing generated an intrathoracic pressure of -10 mmHg and an immediate decrease in ICP by about 7.5 mmHg. The ITD also increased AP (not shown). When the ITD was removed, ICP returned immediately to baseline levels. The impact of both ITD and positive pressure ventilation on ICP suggest a remarkable degree of concordance between changes in intrathoracic and intracranial pressures, which may have significant implications in the treatment of a number of disorders that alter CBF. These new findings also suggest that the vacuum created by the ITD causes a “waterfall” effect that increases blood flow by maximizing the pressure gradient across the cerebral circulation. Maintaining adequate CBF while reducing ICP could prove critical in prolonging or even preventing the progression to circulatory collapse associated with syncope and/or hemorrhagic shock.

Since inadequate cerebral perfusion ultimately leads to syncope and circulatory collapse (Benditt et al., 1996), a device or procedure that effectively maintains or increases CBF might benefit returning astronauts or bleeding patients awaiting definitive medical care. Based on evidence from animal experiments (Lurie et al., 1995, 2002; White et al., 1991) and the observation that subjects reported less severe symptoms (e.g., dizziness) during transition from the squat to standing posture (Convertino et al., 2005b), the effects of ITD breathing on CBF were investigated in humans. Cerebral blood flow velocity (CBFV) was recorded in the right middle cerebral artery in seven subjects using transcranial Doppler ultrasonography. Figure 3 shows a representative response recorded from one subject.




Figure 3. Continuous recording of mean cerebral blood flow (CBF) velocity in a subject before, during (On ITD), and at the cessation (Off ITD) of spontaneous breathing on the ITD.

For all seven subjects, breathing through an ITD increased mean CBFV from 64 cm/s during breathing on a ZTD to 69 cm/s during ITD breathing ($P = 0.01$). End-tidal CO_2 for ITD breathing was 4.8 ± 0.1 %, similar to that produced by the ZTD (4.9 ± 0.2 %) (Cooke et al., 2006). However, it is possible that increased respiratory drive during ITD breathing increased cerebral metabolic activity and therefore induced cerebral vessel dilation. The pulsatility index, an indirect estimate of cerebral vascular resistance tended to decrease with active ITD breathing ($P = 0.09$). The pulsatility index (calculated as the difference between peak systolic and end diastolic flow velocity divided by mean flow velocity) is clearly an imperfect estimate of cerebral vascular resistance that does not take into account systemic arterial, venous, or cerebro-spinal fluid pressures. However, in a prospective study of brain-injured patients, Bellner et al. (2004) found a strong correlation ($r = 0.94$; $p < 0.0001$) between ICP measured by intraventricular catheters and the pulsatility index; they concluded that the latter is a useful surrogate for ICP for monitoring severely brain injured patients (Bellner et al., 2004). Because end-tidal CO_2 is an imperfect predictor of PaCO_2 (van Lieshout et al., 2003), and because even small changes in PaCO_2 profoundly affect CBFV (Ide et al., 2003), it is possible that the observed increases in CBFV during ITD breathing resulted from increased cerebral metabolic activity and consequent dilation of the cerebral vasculature.

6. ORTHOSTATIC STRESS

Resistive breathing might be expected to protect central hemodynamics against circulatory collapse induced by sudden orthostatic stress or hemorrhage (Convertino et al., 2004b, 2004c). One study has addressed this possibility experimentally (Convertino et al., 2005b): Eighteen healthy, normotensive volunteers (9 males, 9 females), ages 20-56, completed two 6-min protocols in counterbalanced order with a ZTD or an ITD set to open at -7 cm H_2O pressure. An infrared finger photoplethysmograph was used to make noninvasive measurements of HR, SV, CO, TPR, and MAP. Symptoms were recorded using a subject perceived rating (SPR) where 1 = normal and 5 = dizziness.

Movement from squat to stand reduced TPR by about 35% with or without the ITD, but the device affected other variables, as illustrated for one subject in Figs. 4 and 5. Using the ZTD, he experienced severe symptoms (SPR = 4) as his SV fell (Fig. 4), AP was reduced and pulse pressure dropped to 40 (Fig. 5). In contrast, the ITD prevented symptoms (SPR = 1), erased the acute, transient drop in SV (Fig. 4) and held pulse pressure at 60 mmHg (Fig. 5). The periodic increases in SV in Fig. 5 reflect the negative intrathoracic pressure induced by the ITD during inspiration.




Fig. 4. Stroke volume responses in a subject undergoing the transition from a squat to a standing posture during spontaneous breathing on an ITD (open circle line) and ZTD (solid line).




Fig. 5. Beat-to-beat arterial blood pressure responses in a subject undergoing the transition from a squat to a standing posture during spontaneous breathing on a ZTD (upper panel) and on an active ITD (lower panel).

On average for all subjects, MAP fell -36 ± 3 mmHg with the ZTD compared to -27 ± 4 mmHg with the ITD ($P = 0.03$) despite similar elevations in HR (15 ± 2 bpm, $P = 0.93$). SV changed by -8 ± 4 % for ZTD vs. $+2 \pm 4$ % for ITD; the corresponding changes in CO were $+10 \pm 6$ % and $+22 \pm 5$ % ($P < 0.04$). The SPR was 1.4 ± 0.1 for ZTD vs. 2.0 ± 0.2 for ITD ($P = 0.04$). These results suggest that the ITD may defend against orthostatic hypotension and intolerance. Future experiments should address the effects of ITD breathing in subjects after a

period of simulated microgravity or experimentally-induced hypovolemia.

7. CENTRAL BLOOD LOSS IN HUMANS

The effects of inspiratory resistance were tested in human volunteers subjected to LBNP as a model for acute reduction of central blood volume due to hemorrhage (Cooke et al., 2004). Figure 6 shows beat-to-beat SV measured with thoracic bioimpedance during baseline supine rest and exposure to 60 mmHg LBNP with normal breathing followed by use of an ITD. In this case, the LBNP caused a 30-35% reduction in SV, while the ITD produced an immediate increase that overshoot and then returned to baseline after 1 min.




Fig. 6. Stroke volume responses in a subject undergoing simulated central blood loss by use of 60 mmHg LBNP during the transition from baseline rest to LBNP to LBNP and spontaneous breathing on an ITD.

Subsequently, we induced central hypovolemia and cardiovascular collapse in nine human volunteers by applying progressive LBNP under counterbalanced experimental conditions with a ZTD or an ITD set to open at -7 cm H₂O pressure. SBP (79 ± 5 mmHg), DBP (57 ± 3 mmHg), and MAP (65 ± 4 mmHg) were lower ($P < 0.02$) when subjects ($n = 9$) breathed through the ZTD than when they breathed through the ITD at the same time of cardiovascular collapse during ZTD breathing (102 ± 3 , 77 ± 3 , 87 ± 3 mmHg). Elevated BP was associated with 23% greater ($P = 0.02$) tolerance to central hypovolemia using an ITD compared with a ZTD (Fig. 7). Thus, in our human model of central blood volume reduction, use of an ITD device increased systemic blood pressure and delayed the onset of cardiovascular collapse during severe hypovolemic hypotension in spontaneously breathing

human volunteers. This device may provide rapid non-invasive hemodynamic support in patients with hypovolemic hypotension once the blood loss has been controlled but before other definitive therapies are available, and therefore provide a critical bridge for maintaining AP in the face of hemorrhage until volume replacement can be provided. Thus, the ITD may prove useful in civilian trauma and tactical combat care, especially for a bleeding patient with a weak or absent pulse.




Fig. 7. LBNP completion (plotted as survival) curves of 9 subjects treated with an ITD (blue curve) and ZTD (red curve).

8. CONCLUSION

Interventions that “buy time” by sustaining adequate perfusion pressures prior to treatment with more definitive therapies are of considerable value for improving patient outcome. In this series of investigations, we demonstrated that improving the efficiency of the intrathoracic vacuum by application of resistance during inspiration with the use of an ITD can increase HR, SV, CO, and arterial blood pressure in humans at rest and during orthostatically-induced central hypovolemia. The elevation in ABP during treatment with an ITD is associated with increased tolerance to severe progressive central hypovolemia leading to cardiovascular collapse and reduced symptoms. Our preliminary results are the first to demonstrate in humans that the time to cardiovascular collapse associated with progressive reduction in central blood volume and subsequent development of severe hypotension can be significantly improved with an ITD. These findings provide compelling evidence that the ITD may extend the therapeutic window of opportunity and improve casualty outcome by allowing greater survival from hemorrhagic shock until definitive care is available. The results of the present investigation suggest that application of an ITD

during the early stages of controlled hemorrhage may restore blood pressure and perfusion to vital organs. While not a panacea, the results from these studies demonstrate that the ITD may help “buy time” until fluid resuscitation is begun, or as an adjunct to conventional resuscitative therapy, thus potentially increasing the likelihood that the casualty can be stabilized and survive to reach a higher echelon of care. It is in this manner that we believe the ITD may provide a critical bridge to more definitive repair of the primary injury and ultimately save lives. In support of this contention, there are currently three confirmed cases from Iraq in which casualties have been successfully treated with the ITD.

ACKNOWLEDGMENTS

The opinions or assertions contained herein are the private views of the authors and are not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense (AR 360-5). The human experimental studies described in this report were reviewed and approved by the Institutional Review Board of the Brooke Army Medical Center, and informed consent was obtained from all participants. The manuscript was reviewed for compliance prior to submission to publication. In conducting the research described here, the authors adhered to the “Guide for the Care and Use of Laboratory Animals”, National Research Council, 1996.

The authors thank Rachel Holder for her assistance in the preparation of this manuscript.

REFERENCES

- Bainbridge, F.A., 1915: The influence of venous filling upon the rate of the heart. *J. Physiol.*, **150**, 65-84.
- Barbieri, R., Triedman, J.K., and Saul, J.P., 2002: Heart rate control and mechanical cardiopulmonary coupling to assess central volume: a systems analysis. *Am. J. Physiol.*, **283**, R1210-R1220.
- Bellner, J., Romner, B., Reinstrup, P., et al., 2004: Transcranial Doppler sonography pulsatility index (PI) reflects intracranial pressure (ICP). *Surg. Neurol.*, **62**, 45-51.
- Benditt, D.G., Lurie, K.G., Adler, S.W., et al., 1996: Pathophysiology of vasovagal syncope. In: Blanc J-J, Benditt D, Sutton R, eds. *Neurally Mediated Syncope: Pathophysiology, Investigations, and Treatment*. Armonk, NY: Futura Publishing; p. 1-24.
- Buckey, J.C., Lane, L.D., Levine, B.D., et al., 1996: Orthostatic intolerance after spaceflight. *J. Appl. Physiol.*, **81**, 7-18.
- Bungo, M.W., Charles, J.B., Johnson, P.C., 1985: Cardiovascular deconditioning during space flight and the use of saline as a countermeasure to orthostatic intolerance. *Aviat. Space Environ. Med.*, **56**, 985-990.
- Carrico, C.J., Holcomb, J.B., Chaudry, I.H., et al., 2002: Post resuscitative and initial utility of life saving efforts. Scientific priorities and strategic planning for resuscitation research and life saving therapy following traumatic injury: report of the PULSE trauma work group. *Acad. Emerg. Med.*, **9**, 621-626.
- Chapleau, M.W., and Abboud, F.M., 1987: Contrasting effects of static and pulsatile pressure on carotid baroreceptor activity in dogs. *Circ. Res.*, **61**, 648-657.
- Chapleau, M.W., and Abboud, F.M., 1989: Determinants of sensitization of carotid baroreceptors by pulsatile pressure in dogs. *Circ. Res.*, **65**, 566-577.
- Convertino, V.A., and Cooke, W.H., 2002: Relationship between stroke volume and sympathetic nerve activity: new insights about autonomic mechanisms of syncope. *J. Gravit. Physiol.*, **9**, P63-P66.
- Convertino, V.A., Cooke, W.H., and Lurie, K.G., 2005: Restoration of central blood volume: application of a simple concept and a simple device to counteract cardiovascular instability in syncope and hemorrhage. *J. Gravit. Physiol.*, **12**, P55-P60.
- Convertino, V.A., Ludwig, D.A., and Cooke, W.H., 2004a: Stroke volume and sympathetic responses to lower-body negative pressure reveal new insight into circulatory shock in humans. *Auton. Neurosci.*, **111**, 127-134.
- Convertino, V.A., Ratliff, D.A., Crissey, J., et al., 2005: Effects of inspiratory impedance on hemodynamic responses to a squat-stand test in human volunteers: implications for treatment of orthostatic hypotension. *Eur. J. Appl. Physiol.*, **94**, 392-399.
- Convertino, V.A., Ratliff, D.A., Ryan, K.L., et al., 2004b: Effects of inspiratory impedance on the carotid-cardiac baroreflex response in humans. *Clin. Auton. Res.*, **14**, 240-248.
- Convertino, V.A., Ratliff, D.A., Ryan, K.L., et al., 2004c: Hemodynamics associated with breathing through an inspiratory impedance threshold device in human volunteers. *Crit. Care Med.*, **32**, S381-S386.
- Convertino, V.A., 1996: Clinical aspects of the control of plasma volume at microgravity and during return to one gravity. *Med. Sci. Sports Exerc.*, **28**, S45-S52.
- Convertino, V.A., Doerr, D.F., Eckberg, D.L., et al., 1990: Head-down bed rest impairs vagal baroreflex responses and provokes orthostatic hypotension. *J. Appl. Physiol.*, **68**, 1458-1464.
- Cooke, W.H., Lurie, K.G., Rohrer, M.J., and Convertino, V.A., 2006: Human autonomic and cerebrovascular responses to inspiratory impedance. *J. Trauma*, **60**, 1275-1283.
- Cooke, W.H., Ryan, K.L., and Convertino, V.A., 2004: Lower body negative pressure as a model to study progression to acute hemorrhagic shock in humans. *J. Appl. Physiol.*, **96**, 1249-1261.

- Cournand, A., Motley, H.L., Werko, L., et al., 1948: Physiological studies of the effects of intermittent positive pressure breathing on cardiac output in man. *Am. J. Physiol.*, **152**, 162-174.
- Eckberg, D.L., and Fritsch, J.M., 1992: Influence of ten-day head-down bedrest on human carotid baroreceptor-cardiac reflex function. *Acta Physiol. Scand.*, **144**, 69-76.
- Eiken, O., Convertino, V.A., Doerr, D.F., Dudley, G.A., et al., 1991: Interaction of the carotid baroreflex, the muscle chemoreflex and the cardiopulmonary baroreflex in man during exercise. *Physiologist*, **34**(Suppl), S118-S120.
- Engelke, K.A., Doerr, D.F., and Convertino, V.A., 1996: Application of acute maximal exercise to protect orthostatic tolerance after simulated microgravity. *Am. J. Physiol.*, **271**, R837-R847.
- Evans, J.M., Stenger, M.B., Moore, F.B., et al., 2004: Centrifuge training increases presyncopal orthostatic tolerance in ambulatory men. *Aviat. Space Environ. Med.*, **75**, 850-858.
- Fitzgerald, R.S., Robotham, J.L., and Anand A., 1981: Baroreceptor output during normal and obstructed breathing and Mueller maneuvers. *Am. J. Physiol.*, **240**, H721-H729.
- Fritsch-Yelle, J.M., Charles, J.B., Jones, M.M., et al., 1994: Spaceflight alters autonomic regulation of arterial pressure. *J. Appl. Physiol.*, **77**, 1776-1783.
- Fritsch, J.M., Charles, J.B., Bennet, B.S., et al., 1992: Short-duration spaceflight impairs human carotid baroreceptor-cardiac reflex responses. *J. Appl. Physiol.*, **73**, 664-671.
- Guyton, A.C., Lindsey, A.W., Abernathy, B., et al., 1957: Venous return at various right atrial pressures and the normal venous return curve. *Am. J. Physiol.*, **189**, 609-615.
- Ide, K., Eliasziw, M., Poulin, M.J., 2003: The relationship between middle cerebral artery blood velocity and end-tidal PCO₂ in the hypocapnic-hypercapnic range in humans. *J. Appl. Physiol.*, **95**, 129-137.
- Kapoor, W.N., 1996: Importance of neurocardiogenic causes in the etiology of syncope. In: Blanc, J.-J., Benditt, D., Sutton, R., eds. *Neurally Mediated Syncope: Pathophysiology, Investigations, and Treatment*. Armonk, NY: Future Publ., p. 55-62.
- Kleiger, R.E., Miller, J.P., Bigger, J.T., and Moss, A.J., 1987: Decreased heart rate variability and its association with increased mortality after acute myocardial infarction. *Am. J. Cardiol.*, **59**, 256-262.
- Levine, B.D., Pawelczyk, J.A., Ertl, A.C., et al., 2002: Human muscle sympathetic neural and haemodynamic responses to tilt following spaceflight. *J. Physiol.*, **538**, 331-340.
- Lurie, K.G., Coffeen, P.R., Shultz, J.J., et al., 1995: Improving active compression-decompression cardiopulmonary resuscitation with an inspiratory impedance valve. *Circulation*, **91**, 1629-1632.
- Lurie, K.G., Voelckel, W., Plaisance, P., et al., 2000: Use of an inspiratory impedance threshold valve during cardiopulmonary resuscitation: a progress report. *Resuscitation*, **44**, 219-230.
- Lurie, K.G., Zielinski, T., McKnite, S., et al., 2002: Use of an inspiratory impedance valve improves neurologically intact survival in a porcine model of ventricular fibrillation. *Circulation*, **105**, 124-129.
- Lurie, K.G., Zielinski, T.M., McKnite, S.H., et al., 2004: Treatment of hypotension in pigs with an inspiratory impedance threshold device: a feasibility study. *Crit. Care Med.*, **32**, 1555-1562.
- Marino, B.S., Yannopoulos, D., Sigurdsson, G., et al., 2004: Spontaneous breathing through an inspiratory impedance threshold device augments cardiac index and stroke volume index in a pediatric porcine model of hemorrhagic hypovolemia. *Crit. Care Med.*, **32**, S398-S405.
- Morgan, B.J., Denahan, T., and Ebert, T.J., 1993: Neurocirculatory consequences of negative intrathoracic pressure vs. asphyxia during voluntary apnea. *J. Appl. Physiol.*, **74**, 2969-2975.
- Pawelczyk, J.A., and Levine, B.D., 1995: Cardiovascular responses to rapid volume infusion: the human Bainbridge reflex. *Circulation*, **92**, I-656.
- Rowell, L.B., and O'Leary, D.S., 1990: Reflex control of the circulation during exercise: chemoreflexes and mechanoreflexes. *J. Appl. Physiol.*, **69**, 407-418.
- Samniah, N., Voelckel, W.G., Zielinski, T.M., et al., 2003: Feasibility and effects of transcutaneous phrenic nerve stimulation combined with an inspiratory impedance threshold in a pig model of hemorrhagic shock. *Crit. Care Med.*, **31**, 1197-1202.
- van Lieshout, J.J., Wieling, W., Karemaker, J.M., et al., 2003: Syncope, cerebral perfusion, and oxygenation. *J. Appl. Physiol.*, **94**, 833-848.
- White, R.J., Leonard, J.I., Srinivasan, R.S., et al., 1991: Mathematical modeling of acute and chronic cardiovascular changes during extended duration orbiter (EDO) flights. *Acta Astronautica*, **23**, 41-51.
- Yannopoulos, D., Sigurdsson, G., McKnite, S., et al., 2005: Effects of incomplete chest wall decompression during cardiopulmonary resuscitation on coronary and cerebral perfusion pressures in a porcine model of cardiac arrest. *Resuscitation*, **64**, 363-372.